

Role of MLL-Wild Type in hematopoiesis and leukemia transformation in MLL-Partial Tandem Duplication mouse models

The James

THE OHIO STATE UNIVERSITY
COMPREHENSIVE CANCER CENTER

Daniel L. Brook¹, Nicholas A. Zorko¹, Adrienne M. Dorrance¹, Daniel A. Yanes¹, Kelsie Bernot¹, Susan P. Whitman¹, Elshafa Ahmed¹, Kathleen McConnell¹, Patricia A. Ernst², Gang Huang³, Guido Marcucci¹, and Michael A. Caligiuri¹
1. Comprehensive Cancer Center, The Ohio State University, Columbus, OH 2. Norris Cotton Cancer Center, Dartmouth College, Hanover, NH 3. Dept. of Experimental Hematology, Cincinnati Children's Hospital, Cincinnati, OH

Background

Acute Myeloid Leukemia (AML) is a blood cancer arising mostly in older adults. MLL-partial tandem duplication (MLL-PTD) is present in 5-7% of cytogenetically normal AML patients. FLT3-internal tandem duplication (FLT3-ITD) is found in 30-35% of AML patients. A subset of AML patients with both FLT3-ITD and MLL-PTD have a poor prognosis.


Single mutant mice expressing either MII-PTD or Flt3-ITD independently do not develop leukemia; however, a double mutant mouse expressing MII-PTD and Flt3-ITD develops AML with a long latency and mimics human AML phenotypically and molecularly¹.

MLL-WT expression is silenced in both human samples² and mouse models³ of MLL-PTD+ AML. Germ-line absence of MII-WT in MII-PTD+ (MII^{PTD/-}) mice is embryonic lethal⁴, so an MII-PTD conditional knockout mouse was obtained to generate MII^{PTD/FLOX};Flt3^{ITD/WT} mice.

Objective 1: Does MII-PTD maintain gain-of-function in the absence of MII-WT?


Objective 2: Is MII-WT a tumor suppressor in an MII-PTD+ AML murine model?

II-PTD;Flt3-ITD cooperate to induce fatal acute leukemia




Survival curve of single knock-in and WT control mice in comparison to the heterozygous double mutant mice.

II^{PTD/WT};Flt3^{ITD/WT} mice with AML develop splenomegaly




Spleens of mice at time of harvest (80 wks).


II^{PTD/WT};Flt3^{ITD/WT} murine acute leukemia is similar to human AML



Objective 1: Does MII-PTD function without MII-WT?




Cre induction deletes MII-FLOX to MII-ΔΔ




The floxed (MII-FLOX) allele is completely deleted, producing MII-ΔΔ in hematopoietic tissues after induction of Cre recombinase.

II-PTD acts as a gain-of-function mutation in the absence of MII-WT



Objective 2: Is MII-WT a tumor suppressor in MII-PTD+ AML?



Experimental design of to test whether MII-WT affects leukemia transformation in MII-PTD+ murine model.

Summary

1. MII-PTD;Flt3-ITD mice develop a transplantable AML that is phenotypically and molecularly similar to human AML.
2. MII^{PTD/ΔΔ} mice maintain normal hematopoiesis survive, but MII^{ΔΔ/ΔΔ} mice succumb to bone marrow failure soon after induction.
3. MII-PTD appears to act as a gain-of-function mutation based on expression levels of the downstream MII targets HoxA9 and Meis1.
4. Breeding of MII^{PTD/FLOX};Flt3^{ITD/WT} with and without Cre mice is in progress to help evaluate the effect of MII-WT on AML transformation.

Future Directions to analyze MII-WT in acute leukemia model

1. Evaluate whether loss of MII-WT (hypothesized to function as a tumor suppressor in MII-PTD+ AML) decreases latency of leukemogenesis in an MII-PTD;Flt3-ITD mouse model of AML.
2. Evaluate expression of oncogenic HoxA9 and Meis1 in MII^{PTD/FLOX};Flt3^{ITD/ITD} mouse model pre- and post- conditional knockout.
3. Monitor disease progression by flow cytometry based immunophenotyping.

References

1. Zorko NA, et al. MII partial tandem duplication and Flt3 internal tandem duplication in a double knock-in mouse recapitulates features of counterpart human acute myeloid leukemias. *Blood*. 2012;120(5):1130-1136.
2. Dorrance, A. M., et al. (2006). MII partial tandem duplication induces aberrant Hox expression in vivo via specific epigenetic alterations. *Journal of Clinical Investigation* 116, 2707-2716.
3. Whitman, S. P., et al. (2005). The MLL partial tandem duplication: evidence for recessive gain-of-function in acute myeloid leukemia identifies a novel patient subgroup for molecular-targeted therapy. *Blood* 106, 345- 352.
4. Jude CD, et al. Unique and independent roles for MLL in adult hematopoietic stem cells and progenitors. *Cell Stem Cell* 1(3);doi:10.1016/ j.stem. 2007.05.019, 2007

Acknowledgements

The following OSUCCC Shared Resources were used to conduct these studies: Nucleic Acid, Flow Cytometry, Comparative Pathology and Mouse Phenotyping. This work was supported by National Cancer Institute grants to MAC (CA089341; CA009338), GM (CA102031), to MAC and GM (CA140158) to NAZ (T32 GM12453, OSU College of Medicine Fellowship and Pelotonia Graduate Fellowship), to KMB (T32 CA009338), and to DB, DY and RS (Pelotonia Undergraduate Fellowship).